



Collaboration for Research, Implementation & Training in Intensive CARE in ASIA

Daily Q Assessment Completed everyday for the inpatient ICU stay. Data reflects events for the previous 24 hrs: 0000-2400. This will include the day of admission.		
Ventilation		Use of mechanical ventilation for any duration in the previous 24 hrs of ICU stay. Nocturnal CPAP from a patient's 'home' system is not considered mechanical ventilation in this context.
Route of mechanical ventilation		If multiple modes please report the most invasive.
Lowest Fio2	% or numerically ≤ 1.0	The lowest recorded inspired oxygen concentration delivered for a minimum of one hour.
Lowest PEEP	mmH2O	The lowest recorded positive end expiratory pressure for a minimum of one hour.
VTE prophylaxis in mechanically ventilated patients		Use of mechanical or pharmacological prophylaxis in the prevention of venous thromboembolism.
VTE prophylaxis Contraindication		The contraindication prevents the use of mechanical and pharmacological prophylaxis.
Stress ulcer prophylaxis in mechanically ventilated patients		Use of proton pump inhibitors or H2 receptor antagonists.
Stress ulcer prophylaxis in mechanically ventilated patients		The named drug from the provided list used for stress ulcer prophylaxis.
Inadvertent/ Accidental extubation		The unexpected or unplanned extubation of a patient.
Highest Temperature	$^{\circ}\text{F}$ or $^{\circ}\text{C}$	Highest recorded temperature.
Cardiovascular Support		Use of continuous intravenous vasoactive medication for a minimum period of one hour.
Vasoactive drugs increased		An increase in the use of vasoactive medication (either number of agents or dosage of an agent) for a minimum period of one hour.
Central venous catheter		The presence of a CVC.
Site		The location of the central venous catheter.
Urinary catheter		The presence of a urinary catheter.
Haemodialysis Catheter		The presence of a haemodialysis catheter.
Site		The location of the haemodialysis catheter.



Collaboration for Research, Implementation & Training in Intensive CARE in ASIA

Renal replacement therapy		Use of renal replacement therapy. Any duration of RRT is considered as yes.
Type of the therapy		The type of renal replacement therapy is either continuous or intermittent.
Sedated		Use of sedative drugs for a minimum of one hour. If bolus sedation, then 2 or more bolus.
Spontaneous breathing trial (SBT)		A trial of spontaneous breathing. An SBT will be defined as one of the four following definitions; <ul style="list-style-type: none"> - Trial of 'T'piece - Pressure support with low support (PS of 5-8) - CPAP (i.e. only PEEP between 0-5) - Automatic tube compensation with CPAP
Highest Wcc	x10 ⁹ /L or x10 ³ /mm ³ or x10 ³ /μL	The highest recorded serum white blood cell count.
Lowest Wcc	x10 ⁹ /L or x10 ³ /mm ³ or x10 ³ /μL	The lowest recorded serum white blood cell count.
Infection		Presence of a confirmed or suspected infection.
Source		The source of infection from an organ list.
Antimicrobial use daily		Use of antimicrobial therapy. Even if only one dose has been administered, this is yes
Antimicrobial type		The type of antimicrobial from a drug list.
Positive Culture		A culture taken which has shown a positive microbial growth.
Date culture taken		The date the sample was taken.
Type of culture		The type of culture taken
Organism		The type of organism found in the culture selected from the drop down list.
Colony forming units (CFU)		For BAL, sputum, Urine cultures: The quantitative measure of the organism found in the culture